

How to rule out cold dark matter

Carlos S. Frenk
Institute for Computational Cosmology,
Durham

The new Ogden
Centre at Durham



Non-baryonic dark matter candidates

From the 1980s:

Type	example	mass
hot	neutrino	few tens of eV
warm	sterile ν	keV-MeV
cold	axion neutralino	$10^{-5}\text{eV} - 100 \text{ GeV}$

The dark matter power spectrum

$k^3 P(k)$

The linear power spectrum (“power per octave”)


Free streaming $\rightarrow$

$\lambda_{\text{cut}} \propto m_x^{-1}$
for thermal relic


$m_{\text{CDM}} \sim 100 \text{ GeV}$
susy; $M_{\text{cut}} \sim 10^{-6} M_{\odot}$

$m_{\text{WDM}} \sim \text{few keV}$
sterile ν ; $M_{\text{cut}} \sim 10^9 M_{\odot}$

$m_{\text{HDM}} \sim \text{few tens eV}$
light ν ; $M_{\text{cut}} \sim 10^{15} M_{\odot}$



Non-linear evolution




Non-linear evolution: simulations

Initial conditions + assumption about content of Universe


Relevant equations:

Collisionless Boltzmann
Poisson, Friedmann eqn
Radiative hydrodynamics
Subgrid astrophysics



How to make a virtual universe

Non-baryonic dark matter cosmologies




Neutrinos
 $\Omega=1$

Frenk, White
& Davis '83

Non-baryonic dark matter cosmologies

Neutrino DM →
wrong clustering

Neutrinos cannot
make appreciable
contribution to Ω
→ $m_\nu \ll 30$ eV




Non-baryonic dark matter cosmologies

Neutrino DM →
wrong clustering

Neutrinos cannot
make appreciable
contribution to Ω
→ $m_\nu \ll 30$ eV



Early CDM N-body
simulations gave
promising results

In CDM structure
forms hierarchically




The Λ CDM model of cosmogony

Proposed in 1980s Now empirically supported by:



Springel, Frenk &
White 2006

The cosmic power spectrum: from the CMB to the 2dFGRS




$z \sim 1000$

$\text{Log } k^3 P(k)$

wavelength k^{-1} (comoving h^{-1} Mpc)


1 000 100 10



$z \sim 0$

⇒ Λ CDM provides an excellent description of mass power spectrum from 10-1000 Mpc

Sanchez et al 06



The cosmic power spectrum: from the CMB to the 2dFGRS

Free streaming $\rightarrow$

$$\lambda_{\text{cut}} \propto m_x^{-1}$$

for thermal relic


$$m_{\text{CDM}} \sim 100 \text{ GeV}$$

$$\text{susy}; M_{\text{cut}} \sim 10^{-6} M_{\odot}$$

$$m_{\text{WDM}} \sim \text{few keV}$$

$$\text{sterile } \nu; M_{\text{cut}} \sim 10^9 M_{\odot}$$

Log $k^3 P(k)$ wavelength k^{-1} (comoving h^{-1} Mpc)





Both CDM & WDM compatible with CMB & galaxy clustering

Claims that both types of DM have been discovered:

- ◆ CDM: γ -ray excess from Galactic Center
- ◆ WDM (sterile ν): 3.5 X-ray keV line in galaxies and clusters

Very unlikely that both are right!

The cosmic power spectrum: from the CMB to the 2dFGRS

Free streaming $\rightarrow$

$$\lambda_{\text{cut}} \propto m_x^{-1}$$

for thermal relic


$$m_{\text{CDM}} \sim 100 \text{ GeV}$$

$$\text{susy}; M_{\text{cut}} \sim 10^{-6} M_{\odot}$$

$$m_{\text{WDM}} \sim \text{few keV}$$

$$\text{sterile } \nu; M_{\text{cut}} \sim 10^9 M_{\odot}$$

Log $k^3 P(k)$ wavelength k^{-1} (comoving $h^{-1} \text{ Mpc}$)



Sterile neutrinos

Explain:

- Neutrino oscillations and masses
- Baryogenesis
- Absence of right-handed neutrinos in standard model
- Dark matter

Sterile neutrino minimal standard model (ν MSM; Boyarski+ 09):


- Extension of SM w. 3 sterile neutrinos: 2 of GeV; 1 of keV mass
- If $\Omega_N = \Omega_{DM}$, 2 parameters: mass, lepton asymmetry/mixing angle
- GeV particles may be detected at CERN (SHiP)
- Dark matter candidate can be detected by X-ray decay

Primordial $P(k)$ for 7 keV sterile neutrino models

- Thermal and resonant production mechanisms
- Resonant production depends on baryon asymmetry parameter, L_6
- Linear PS varies **non-monotonically** with L_6

Ly- α forest rules out thermal masses, $m_\nu < 3.3$ keV (Viel + '13)

Lovell, Bose, CSF et al. 16





Cold Dark Matter

Warm Dark Matter

13.4 billion years ago

cold dark matter

warm dark matter

How can we distinguish between these?

Lovell, Eke, Frenk, Gao, Jenkins, Wang, White, Theuns,
Boyarski & Ruchayskiy '12


cold dark matter

warm dark matter

Obvious to

In the M

Th



MW or M31

ered so far

G!

Lovell, Eke, Frenk, Gao, Jenkins, Wang, White, Theuns,
Boyarski & Ruchayskiy '12


Most subhalos never make a galaxy!

Because:

- Reionization heats gas to 10^4K , preventing it from cooling and forming stars in small halos ($T_{\text{vir}} < 10^4\text{K}$)
- Supernovae feedback expels residual gas in slightly larger halos


Luminosity Function of Local Group Satellites

- Median model → correct abund. of sats brighter than $M_V = -9$ and $V_{\text{cir}} > 12$ km/s
- Model predicts many, as yet undiscovered, faint satellites
- LMC/SMC should be rare (~10% of cases)



Luminosity Function of Local Group Satellites

- Median model → correct abund. of sats brighter than $M_V = -9$ and $V_{\text{cir}} > 12$ km/s
- Model predicts many, as yet undiscovered, faint satellites
- LMC/SMC should be rare (~10% of cases)



Benson, Frenk, Lacey, Baugh & Cole '02
(see also Kauffman+ '93, Bullock+ '00, Somerville '02)



VIRGO

icc.dur.ac.uk/Eagle

“Evolution and assembly of galaxies and
their environment”

THE EAGLE PROJECT

Virgo Consortium


Durham: Richard Bower, Michelle Furlong, Carlos Frenk, Matthieu Schaller, James Trayford, Yelti Rosas-Guevara, Tom Theuns, Yan Qu, John Helly, Adrian Jenkins.

Leiden: Rob Crain, Joop Schaye.

Other: Claudio Dalla Vecchia, Ian McCarthy, Craig Booth...

Galaxy stellar mass function

Comparison to semi-analytic models



VIRG

Dark matter

APOSTLE
EAGLE full
hydro
simulations

Local Group

CDM

Sawala et al '16



Stars

VIRG

APOSTLE
EAGLE full
hydro
simulations

Local Group


Stars

Far fewer satellite galaxies than CDM halos

Sawala et al '16



EAGLE Local Group simulation






The Auriga MW-like galaxies

Grand et al '16

30 very high res
Aureo sims


6 even higher
res sims

D. Campbell
C. Frenk
F. Gomez
R. Grand
A. Jenkins
F. Marinacci
R. Pakmor
V. Springel
S. White



Fraction of dark subhalos

$$V_c = \sqrt{\frac{GM}{r}} \quad V_{\max} = \max V_c$$



All halos of mass $< 5 \times 10^8 M_{\odot}$ or $V_{\max} < 7 \text{ km/s}$ are dark ($m_* < 10^4 M_{\odot}$)



(~50 discovered so far)



(a few tens)



Warm DM: different ν mass

$z=3$

WDM

2.3 keV

2.0 keV

1.6 keV

1.4 keV

CDM






WDM

1.4keV

2.3keV


2.0keV

1.6keV



Luminosity Function of Local Group Satellites in WDM

From “Warm Apostle:” 7keV sterile ν $M_h \sim 10^{12} M_\odot$



Lovell et al. '16

When “baryon effects” are
taken into account



Observed abundance of satellites
is compatible with CDM but rules
out some WDM models



There is no such thing as the
“satellite problem” in CDM!




All we have achieved by
counting satellite galaxies
is to rule out a few WDM
models!

Does the inner
structure of satellites
help?



The Density Profile of Cold Dark Matter Halos



Shape of halo profiles
~independent of halo mass &
cosmological parameters

Density profiles are “cuspy” -
no ‘core’ near the centre

Fitted by simple formula:

$$\frac{\rho(r)}{\rho_{crit}} = \frac{\delta_c}{(r/r_s)(1+r/r_s)^2}$$

(Navarro, Frenk & White '97)

More massive halos and
halos that form earlier have
higher densities (bigger δ)



The core-cusp problem

cold dark matter

warm dark matter

Halos and subhalos in CDM & WDM have
cuspy NFW profiles

$$\frac{\rho(r)}{\rho_{crit}} = \frac{\delta_c}{(r / r_s)(1 + r / r_s)^2}$$

Lovell, Eke, Frenk, Gao, Jenkins, Theuns '12



How about baryon effects?

The cores of dwarf galaxy haloes

Julio F. Navarro,^{1,2★} Vincent R. Eke² and Carlos S. Frenk²

¹*Steward Observatory, The University of Arizona, Tucson, AZ 85721, USA*

²*Physics Department, University of Durham, South Road, Durham DH1 3LE*

Accepted 1996 September 2. Received 1996 August 28; in original form 1996 June 26

ABSTRACT

We use N -body simulations to examine the effects of mass outflows on the density profiles of cold dark matter (CDM) haloes surrounding dwarf galaxies. In particular, we investigate the consequences of supernova-driven winds that expel a large fraction of the baryonic component from a dwarf galaxy disc after a vigorous episode of star formation. We show that this sudden loss of mass leads to the formation of a core in the dark matter density profile, although the original halo is modelled by a coreless (Hernquist) profile. The core radius thus created is a sensitive function of the mass and radius of the baryonic disc being blown up. The loss of a disc with mass and size consistent with primordial nucleosynthesis constraints and angular momentum considerations imprints a core radius that is only a small fraction of the original scalelength of the halo. These small perturbations are, however, enough to reconcile the rotation curves of dwarf irregulars with the density profiles of haloes formed in the standard CDM scenario.

Let gas cool and condense to the galactic centre

→ gas self-gravitating
→ star formation/burst

Rapid ejection of gas during starburst → a core in the halo dark matter density profile

Navarro, Eke, Frenk '96

Parry, CSF et al. '11

Governato et al. '12

Pontzen & Governato '12

Brooks et al. '12

Navarro, Eke, Frenk '96

The cores of dwarf galaxy haloes L75





Figure 3. Equilibrium density profiles of haloes after removal of the disc. The solid line is the original Hernquist profile, common to all cases. The dot-dashed line is the equilibrium profile of the 10 000-particle realization of the Hernquist model run in isolation at $t=200$. (a) $M_{\text{disc}}=0.2$. (b) $M_{\text{disc}}=0.1$. (c) $M_{\text{disc}}=0.05$.

Cores in dwarf galaxy simulations

Governato et al. assume
high density threshold for
star formation

EAGLE does not

- High threshold allows
large gas mass to
accumulate in centre
- Sudden repeated
removal of gas transfers
binding energy




Governato et al. '12

Pontzen et al. '12



Cores in dwarf galaxy simulations

Cores may only form in a limited range of galaxy masses

- Galaxy too small → not enough energy
- Galaxy too large → potential well too deep





Di Cintio et al. '14



Dwarf galaxies in Apostle have NFW cusps!

Sawala et al '15



Does Nature have them?



Sawala et al '15



Dwarf galaxies around the Milky Way

Fornax

Sculptor

Leo I

© Anglo-Australian Observatory

Carina

Sextans

Sagittarius

The DM halo of the Sculptor dwarf

Sculptor has two stellar pops:


- (i) centrally concentrated, high [Fe/H]
- (ii) extended, low [Fe/H]

$$M(< r) = \mu \frac{r < \sigma_{los}^2 >}{G}$$

↑
 $r=r_{1/2}$

Walker '10; Wolf et al '10 →

if $r=r_{1/2}$, $\mu=2.5$, independently of model assumptions!




The DM halo of the Sculptor dwarf

$$M(< r) = \mu \frac{r \langle \sigma_{los}^2 \rangle}{G}$$

Key assumption of mass estimator:

spherical symmetry

Is Sculptor spherical?



Genina, CSF et al '17

The DM halo of the Sculptor dwarf


$$M(< r) = \mu \frac{r < \sigma_{los}^2 >}{G}$$

Key assumption of mass estimator:

spherical symmetry

Most satellites in apostle are elongated!

View galaxy from different directions



You can infer any slope, from NFW to core depending on viewing angle!

Genina, CSF et al '17



Many nearby galaxies now have hi-res 2D HI velocity fields → ideal for inferring potential

Assume: gas is in centrifugal equilibrium on approximately circular orbits

Rotation curves of 2 APOSTLE dwarfs

APOSTLE galaxies all have NFW cusps

2D velocity field $\rightarrow V_c(r)$ (rotn curve); in dynamical equilibrium: $V_c = \sqrt{\frac{GM(<r)}{r}}$



Rotation curves of 2 APOSTLE dwarfs

APOSTLE galaxies all have NFW cusps

2D velocity field $\rightarrow V_c(r)$ (rotn curve); in dynamical equilibrium: $V_c = \sqrt{\frac{GM(<r)}{v}}$

$\downarrow$

$\bullet$ 3D BAROLO fit

Tilted-ring model corrected for asymmetric drift



Let's apply this to APOSTLE galaxies by making a mock 2D velocity field and analysing it just as the real data

Rotation curves of 2 APOSTLE dwarfs

APOSTLE galaxies all have NFW cusps

2D velocity field $\rightarrow V_c(r)$ (rotn curve); in dynamical equilibrium: $V_c = \sqrt{\frac{GM(<r)}{v}}$

$\downarrow$ Tilted-ring model corrected for asymmetric drift
 $\bullet$ ${}^{\text{3D}}$ BAROLO fit



Oman et al '17; Marasco et al '17

Conclusions from halo structure

No evidence for cores in dwarf galaxies → inferred 'cores' due to systematics in analysis of stellar dynamics and HI rotation curves:

If cores exist → can be produced by baryon effects



There is NO evidence for
cores in dwarf galaxies

Existing data are consistent
with either cusps or cores





So, we can't distinguish
CDM from WDM by
counting satellite galaxies

There is no need for
despair: there is a way
to distinguish them





Can we distinguish CDM/WDM?

cold dark matter

warm dark matter

Rather than counting faint galaxies,
count the number of dark halos

The subhalo mass function




CDM

WDM

3 x fewer WDM subhalos at $3 \times 10^9 M_\odot$

10 x fewer at $10^8 M_\odot$





Can we distinguish CDM/WDM?

cold dark matter

warm dark matter


1. Gaps in stellar streams (PAndAS, GAIA)
2. Gravitational lensing




Can we distinguish CDM/WDM?

Cooper et al '16

CDM simulation



IC1102



Subhalos crossing a cold tidal stream can produce a gap

Globular cluster streams (e.g. Pal 5) may be best



Can we distinguish CDM/WDM?

cold dark matter

warm dark matter


1. Gaps in stellar streams (PAndAS, GAIA)
2. Gravitational lensing



Gravitational lensing: Einstein rings

How to rule out CDM

Gravitational lensing: Einstein rings




When the source and the lens are well aligned → strong arc or an Einstein ring



SLAC sample of strong lenses

Einstein Ring Gravitational Lenses


Hubble Space Telescope • ACS



NASA, ESA, A. Bolton (Harvard-Smithsonian CfA), and the SLACS Team

STScI-PRC05-32

Gravitational lensing: Einstein rings



When the source and the lens are well aligned → strong arc or an Einstein ring



Gravitational lensing: Einstein rings


Halos projected onto an Einstein ring distort the image



Vegetti & Koopmans '09

Detecting substructures with strong lensing

Vegetti & Koopmans '09





Can detect subhalos as small as $10^7 M_{\odot}$

Substructures vs interlopers

Subhalos & halos projected along the l.o.s both lens: who wins?

Projected l.o.s halos




The number of line-of-sight haloes is larger than that of subhaloes


Substructures vs interlopers

Subhalos & halos projected along the l.o.s both lens: who wins?

Projected l.o.s halos



Li, CSF et al. '16



→ This is the **cleanest** possible **test**: it depends **ONLY** on the **small-mass** end of the “**field**” halo mass function which we know how to calculate and is **unaffected by baryons**


Detecting substructures with strong lensing

Σ_{tot} = projected halo number density within Einstein ring

m_c = halo cutoff mass

100 Einstein ring systems and detection limit: $m_{\text{low}} = 10^7 h^{-1} M_\odot$

Detection limit = $10^7 h^{-1} M_\odot$



m_c = halo cutoff mass

$m_c = 1.3 \times 10^8 h^{-1} M_\odot$ for coldest 7 keV sterile neutrino

- If DM is 7 keV sterile $\nu \rightarrow$ **exclude** CDM at $\gg \sigma$!
- If DM is CDM $\rightarrow$ **exclude** 7 keV sterile ν at $\gg \sigma$



Conclusions

- Λ CDM: great **success** on scales $> 1\text{Mpc}$: CMB, LSS, gal evolution
 - But on these scales **Λ CDM** cannot be distinguished from **WDM**
 - The **identity** of the DM makes a big difference on **small scales**
1. Counting faint galaxies **cannot** distinguish CDM/WDM
 2. Halos $< \sim 5 \cdot 10^8 M_0$ are dark; halos $> 10^{10} M_0$ are bright
(abundance matching fails for halos $< 10^{10} M_0$)
 3. No evidence for cores but baryon effects can make them
 4. Distortions of **strong** gravitational **lenses** offer a **clean**
test of CDM vs WDM $\rightarrow$ and can potentially **rule out** CDM!